

**Supplementary Material 2. ICD-10 codes for comorbidities**

<b>Comorbidities</b>	<b>ICD-10 codes</b>
<b>Hypertension</b>	I10 (essential hypertension), I13 (hypertensive heart and renal disease), or I15 (secondary hypertension)
<b>Diabetes</b>	E11 (non-insulin-dependent diabetes mellitus), E12 (malnutrition-related diabetes mellitus), E13 (other specified diabetes mellitus), or E14 (unspecified diabetes mellitus)
<b>Dyslipidemia</b>	E78 (disorders of lipoprotein metabolism and other lipidemias)
<b>Myocardial infarction</b>	I21 (ST elevation and non-ST elevation myocardial infarction), I22 (subsequent ST elevation and non-ST elevation myocardial infarction), I23 (current complications following ST elevation and non-ST elevation myocardial infarction), I24 (other acute ischemic heart diseases), or I25 (chronic ischemic heart disease)
<b>Congestive heart failure</b>	I50.0 (congestive heart failure)
<b>Stroke</b>	I63 (cerebral infarction) or I64 (stroke, not specified as hemorrhage or infarction)
<b>Depression</b>	F32 (depressive episode) or F33 (recurrent depressive disorder)
<b>Anxiety</b>	F40 (phobic anxiety disorders) or F41 (other anxiety disorders)

Abbreviation: ICD-10, International Classification of Diseases, Tenth Revision